

POSTER PRESENTATION

Open Access

Pancreatic cancers depend on a non-canonical glutamine metabolism pathway

Costas Lyssiotis^{1*}, Jaekyoung Son², Joesph Mancias², Haoqiang Ying³, Alec Kimmelman², Lewis Cantley¹

From *Metabolism, Diet and Disease 2014: Cancer and metabolism*
Washington DC, USA. 28-30 May 2014

Cancer cells exhibit metabolic dependencies that distinguish them from their normal counterparts. Among these addictions is an increased utilization of the amino acid glutamine (Gln) to fuel anabolic processes. Recently, we reported the identification of a non-canonical pathway of Gln utilization in human pancreatic cancer cells that is required for tumor growth. While most cells utilize glutamate dehydrogenase (GLUD1) to convert Gln-derived glutamate (Glu) into α -ketoglutarate (α KG) in the mitochondria to fuel the tricarboxylic acid cycle, pancreatic cancer cells rely on a distinct pathway that integrates the mitochondrial and cytosolic aspartate aminotransferases GOT2 and GOT1. By generating α KG from Glu (in conjunction with the conversion of oxaloacetate into aspartate), GOT2 fuels anaplerosis in place of GLUD1. The Asp created is released into the cytosol and acted on by GOT1. This is subsequently used through a series of reactions to yield cytosolic NADPH from malic enzyme. Importantly, we have demonstrated that pancreatic cancers are strongly dependent on this series of reactions to maintain redox homeostasis which enables proliferation. Herein, we detail the subcellular compartmentalization and consequences of the aforementioned reactions on pancreatic cancer metabolism. We have also investigated the essentiality of this pathway in other contexts and find that pancreatic cancers have a uniform and unique reliance on this pathway, which may provide novel therapeutic approaches to treat these refractory tumors.

Authors' details

¹Weill Cornell Medical College, New York, NY, USA. ²Dana-Farber Cancer Institute, Boston, MA, USA. ³MD Anderson Cancer Center, Houston, TX, USA.

Published: 28 May 2014

doi:10.1186/2049-3002-2-S1-P44

Cite this article as: Lyssiotis et al.: Pancreatic cancers depend on a non-canonical glutamine metabolism pathway. *Cancer & Metabolism* 2014 **2**(Suppl 1):P44.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Weill Cornell Medical College, New York, NY, USA
Full list of author information is available at the end of the article